

Investigating the Effect of Quinine-Induced Autophagy Inhibition on the Efficacy of Curcumin Cancer Treatment

Ezeck Chong (4S2), Yeo Rong Quan (4S2)

Group 1-20

Abstract

Overexposure to UVB radiation in sunlight can cause DNA damage and increase the risk of certain types of cancer. Autophagy is a normal physiology process where dysfunctional or damaged cellular contents are degraded and recycled during cellular stress. In the early stages of tumour development, autophagy plays a pro death role by removing damaged organelles and degrading tumour-protecting proteins. In the late stages of tumour development, its pro-survival role helps cancer cells resist cytotoxic stress from anti-cancer drugs. This study aims to investigate the optimal concentration of quinine, a known an autophagy inhibitor, with curcumin and goji berry in restoring the maximum survival of UV-treated *Caenorhabditis elegans*, a good model organism for anti-cancer drug discovery and cancer research. Curcumin and goji berry were selected as they are known to possess anticancer and antioxidant properties. It was found that the survival rates for UV-treated *C. elegans* added with curcumin had a significant increase as compared to those not added with curcumin. When different concentrations of quinine were added with curcumin, the survival rates did not increase. However, the survival rates for UV-treated *C. elegans* with goji berry showed a significant increase as compared to those without this compound. Different concentrations of quinine with goji berry promoted higher survival rates than those with goji berry alone. The optimal concentrations of quinine that restored the highest survival of *C. elegans* were found to be 80 µg/ml and 60 µg/ml when used in combination with curcumin and goji berry, respectively. These results can be applied in the treatment of initial stages of DNA damage induced by UV radiation, hence can potentially be used as effective anticancer agents.

Introduction

Exposure to sunlight for prolonged periods can damage the skin, cause sunburn and increase the risk of certain types of cancer. The medium-wavelength UVB component of sunlight is one of the major risk factors for the development of skin cancer in humans. UVB light is known to induce oxidative stress by generating reactive oxygen species (ROS), which are responsible for oxidative damage to proteins, DNA, RNA and lipids (Perluigi et al., 2010).

Autophagy is a normal physiology process where dysfunctional or damaged cellular contents are degraded and recycled during cellular stress. Autophagy plays an important role in cancer because of its tumour inhibiting and tumour protecting functions. In the early stages of tumour development, autophagy helps healthy cells suppress tumour growth by maintaining cellular homeostasis and degrading tumour-promoting proteins. However, in the later stages, autophagy helps the cancer cells resist cytotoxic stress from anti-cancer drugs. Quinine, which is a quinoline-based antimalarial compound, is a known autophagy inhibitor (Golden et al., 2015).

The genome of the nematode *Caenorhabditis elegans* has a high similarity with the human genome, hence making it a good model organism for anti-cancer drug discovery and cancer research (Kobet et al., 2014; Kyriakakis, Markaki, & Tavernarakis, 2014). UV-B light (302 nm) is used as a convenient way to induce large amounts of DNA damage in *C. elegans* (Lans & Vermeulen, 2011). This may model early stage cancer because of DNA damage.

Curcumin, a polyphenolic compound from the turmeric root has been researched extensively for the treatment of cancer in animals because of its anticancer and antiproliferative properties. It can inhibit tumor initiation and tumor promotion with minimal side effects (Ravindran, Prasad, & Aggarwal, 2009; Vallianou, Evangelopoulos, Schizas, & Kazazis, 2015; Zhou et al., 2017). Goji berry, scientifically known as *Lycium barbarum*, has been used in Traditional Chinese Medicine for more than 2000 years. It has been studied as a potential anticancer agent in humans because of its antioxidant and antiproliferative properties (Zhou et al., 2014; Wawruszak, Czerwonka, Okła, & Rzeski, W., 2015).

Few studies have combined both curcumin and quinine in the treatment of cancer. This study aimed to investigate if inhibiting autophagy using quinine enhanced the anticancer effects of curcumin and goji berry. The effects of curcumin and goji berry added with various concentrations of quinine on UVB-treated *C. elegans* were studied.

Objectives and Hypothesis

The objective of this study was to investigate the optimal concentration of quinine an autophagy inhibitor with curcumin and goji berry in restoring the maximum survival of UV-treated *C. elegans*. We hypothesise that curcumin and goji berry could protect *C. elegans* against UV light-induced DNA damage, restoring survival and locomotion of *C. elegans*. Greater concentration of quinine, an autophagy inhibitor, with curcumin or goji berry would lead to decreased protection of *C. elegans* against UV-induced oxidative stress.

Experimental Procedures

Preparation of curcumin, goji berry and quinine solutions

0.01g of curcumin was dissolved in 100ml of water and it was sterilized with a sterilized filter. 0.01g of quinine was dissolved in 100ml of water and it was sterilized with a sterilized filter. 10g of goji berries were ground with a mortar and pestle and mixed with 100ml of water. The mixture was centrifuged for 10min at 7000 rpm and sterilized with a sterilized filter.

DPPH test

The DPPH test was conducted to determine the antioxidant activity of curcumin and goji berry. 1,1-diphenyl-2-picryl-hydrazil (DPPH) is a free radical which produces a purple solution when dissolved in methanol. When it is reduced by antioxidants, a change in colouration from purple to yellow is observed. The negative control consisted of 1.0 ml of DPPH, 1.9 ml of methanol and 0.1 ml of sterile water. In test set-ups, curcumin and goji berry replaced the sterile water. Five replicates were prepared. For the respective blanks for each set-up, methanol was added instead of DPPH solution. The initial absorbance was measured at 517 nm against the respective blanks and the mixtures were then left to stand in the darkness for 20 min, before the final absorbance readings were measured.

The % radical scavenging activity was calculated as follows:

$(\text{Final absorbance of control} - \text{Final absorbance of test}) / \text{Final absorbance of control} \times 100\%$

Growth of bacteria

Escherichia coli OP50 was inoculated and grown in 10 ml of LB broth overnight at 30°C in a shaking incubator.

Preparation of NGM medium for *C. elegans*

The composition of NGM is as follows: 0.9 g NaCl, 7.5 g agar, 0.75 g bacto peptone in 300 ml water. After autoclaving, 0.3 ml cholesterol (5 mg/ml), 0.3 ml MgSO₄ (1 M), 0.3 ml CaCl₂ (1 M), 7.5 ml potassium phosphate buffer pH 6.0 (1 M) are added.

Treatment of *C. elegans* with curcumin, goji berry and quinine

Curcumin, goji berry and quinine were dissolved in water and 0.5 ml of each were spread onto NGM plates. 0.05 ml of *E. coli* OP50 broth culture were added to these NGM plates and grown overnight at 30°C. A block of agar containing *C. elegans* N2 was placed at the centre of the NGM plate. The plate was incubated at 20°C for 2 days for the growth of *C. elegans*. After 2 days, *C. elegans* were collected with M9 buffer and centrifuged for 1 min at 2000 rpm.

Inducing oxidative stress in *C. elegans*

A Petri dish containing the mixtures in the test and control setups was placed on the UV transilluminator. UV light of wavelength of 302 nm was shone on *C. elegans* for 2 min to induce DMA damage and oxidative stress. Five replicates were prepared. The percentage of *C. elegans* that survived was determined. A summary of the setups are presented in Table 1.1.

Table 1.1: Summary of treatments

Test	UV-treated <i>C. elegans</i> treated with curcumin/goji berry and quinine
Control (1)	UV-treated <i>C. elegans</i> treated with curcumin/goji berry only
Control (2)	UV-treated <i>C. elegans</i> without curcumin/goji berry
Control (3)	No UV-treated <i>C. elegans</i>

Data analysis

The Kruskal-Wallis test was carried out to determine if there are significant differences in the mean percentage survival of *C. elegans* among the setups.

Results and Discussions

Goji berry was shown to be antioxidant through the DPPH test with a radical scavenging activity (RSA) of 38.6%. There was no significant difference of absorbance between the control and the curcumin setup. (Mann-Whitney U test p value of 0.144). However there was a significant difference of absorbance between the control and the goji berry setup (Mann-Whitney U test p value of 0.012)




Fig. 1.1: DPPH antioxidant test for curcumin and goji berry extracts

UV-treated *C. elegans* added with curcumin resulted in a significant increase in the percentage survival as compared to those without addition of curcumin (Mann-Whitney U test p value of 0.012 for setups UV vs UV+C.) There was a significant difference between the survival percentage of curcumin and curcumin with quinine setups (Kruskal Wallis test p value of 0.000). As the concentration of quinine increased, the percentage of survived UV-treated *C. elegans* increased, although it was lower than treatment with curcumin alone. The setup with the highest percentage of UV-treated *C. elegans* that survived with curcumin and quinine treatment occurred when the quinine concentration was 80 µg/ml. However, there was no significant difference between the percentage survival of UV-treated *C. elegans* with curcumin alone and this particular setup (Kruskal Wallis test p value of 0.117 for setups UV+C vs UV+C+Q80). C represents curcumin only and C+Q80 represents the setup with curcumin and 80 µg/ml quinine. Results are shown in Fig. 1.2.




Fig. 1.2: Effect of curcumin and quinine on survival of UV-treated *C. elegans*

UV-treated *C. elegans* added with curcumin resulted in a significantly higher speed of movement as compared to those without curcumin (Mann-Whitney U test p value of 0.000 for setups UV vs UV+C.) There was a significant difference between the speed of movement of curcumin and curcumin with quinine setups (Kruskal Wallis test p value of 0.000). As the concentration of quinine increased, the speed of movement of UV-treated *C. elegans* increased, although it was still slower than treatment with curcumin alone. The setup with the highest speed of movement of curcumin and quinine treatment occurred when the quinine concentration was at 80 $\mu\text{g/ml}$. However, there was no significant difference between the speed of movement of UV-treated *C. elegans* with curcumin alone and this particular setup (Kruskal Wallis test p value of 0.321 for setups UV+C vs UV+C+Q80). Results are shown in Fig. 1.3.




Fig. 1.3: Effect of curcumin and quinine on the locomotion of UV-treated *C. elegans*

UV-treated *C. elegans* added with goji berry extract resulted in a significant increase in percentage survival as compared to those not treated with goji berry (Mann-Whitney U test p value of 0.012 for setups UV vs UV+G). There was a significant difference between the survival percentage of goji berry alone and goji berry with quinine setups (Kruskal Wallis test p value of 0.001). As the concentration of quinine increased, the percentage of UV-treated *C. elegans* that survived increased. The optimal treatment with the highest percentage of survived UV-treated *C. elegans* was with goji berry and quinine at concentrations of 60 µg/ml, i.e. setup UV+G+Q60. The results are shown in Fig. 1.4.




Fig. 1.4: Effect of goji berry and quinine on survival of UV-treated *C. elegans*

UV-treated *C. elegans* fed with goji berry resulted in a significantly higher speed of movement as compared to those without goji berry treatment (Mann-Whitney U test p value of 0.000 for setups UV vs UV+G). There was a significant difference between the speed of movement of goji berry and goji berry with quinine setups (Kruskal Wallis test p value of 0.000). As the concentration of quinine increased, the speed of movement of UV-treated *C. elegans* increased. The optimal treatment with the highest speed of movement was with goji berry and quinine at concentrations of 60 µg/ml, i.e. setup UV+G+Q60. Results are shown in Fig. 1.5.




Fig. 1.5: Effect of goji berry and quinine on the locomotion of UV-treated *C. elegans*

Autophagy plays both a pro-survival and pro-death role in cells. When subjected to stresses like starvation, autophagy remove damaged organelles or protein aggregates and recycle metabolic substrates to maintain energy homeostasis, playing a pro-survival role. However, excessive levels of autophagy promote autophagic cell death and plays a pro-death role (Kang, You & Avery, 2007; Dalby, Tekedereli, Lopez-Berestein, & Ozpolat, 2010).

The results of this study have shown that quinine, a known autophagy inhibitor, has different effects when added to curcumin and goji berry. Curcumin is a naturally occurring autophagy modulator that can remove dysfunctional organelles and provide an energy source or essential nutrients to maintain cellular activity (Ravindran et al., 2009; Shakeri, Cicero, Panahi, Mohajeri & Sahebkar, 2018). Addition of quinine seemed to interfere with the pro-death role of curcumin to remove damaged organelles and made it less effective in promoting the survival of *C. elegans*. Goji berry has been reported to have anti-oxidant and anti-cancer properties (Zhou, et al., 2014; Wawruszak et al., 2015). Addition of quinine to goji berry treatment increased the survival rate of UV-treated *C. elegans* as it enhanced its pro-survival role. Therefore, different levels of autophagy when used in combination with curcumin or goji berry either increased or decreased the survival of *C. elegans*.




Fig. 1.6: UV-treated *C. elegans* without curcumin or goji berry




Fig. 1.7: UV-treated *C. elegans* with goji berry

UV-treated *C. elegans* treated without curcumin or goji berry looked non motile and the body was straight suggesting that they were dead. UV-treated *C. elegans* treated with goji berry looked motile and the body was curled up suggesting that they were alive.

Conclusions

The results of this study showed that the optimal concentrations of quinine, an autophagy inhibitor, that restored the highest survival of *C. elegans* were found to be 80 µg/ml and 60 µg/ml when used in combination with curcumin and goji berry, respectively. In the initial stages of DNA damage, quinine should not be added to curcumin as this reduced the efficacy of curcumin to remove damaged organelles. In contrast, the addition of quinine with goji berry helped in the DNA damage, increasing the survival of UV-treated *C. elegans*.

A limitation of this study is that UVB radiation causes DNA damage which is a precursor to cancer, but fluorescence microscopy analysis needs to be performed to establish the level of autophagy under the various treatments. Despite the fact that *C. elegans* is a good model organism to show human diseases, it is not known if the same trend observed in this study can be extended to human cancer cell lines.

As future work, the level of autophagy can be monitored using fluorescence microscopy so that the effect of quinine can be better ascertained. A combination of curcumin and goji berry can also be tested to determine if any synergistic effects in restoring survival of UV-exposed *C. elegans* can be observed.

References

- Dalby, K., Tekedereli, I., Lopez-Berestein, G., & Ozpolat, B. (2010). Targeting the pro-death and pro-survival functions of autophagy as novel therapeutic strategies in cancer. *Autophagy*, 6(3), 322-329. doi:10.4161/auto.6.3.11625
- Golden, E. B., Cho, H., Hofman, F. M., Louie, S. G., Schönthal, A. H., & Chen, T. C. (2015). Quinoline-based antimalarial drugs: A novel class of autophagy inhibitors. *Neurosurgical Focus*, 38(3), E12. doi:10.3171/2014.12.focus14748
- Kang, C., You, Y., & Avery, L. (2007). Dual roles of autophagy in the survival of *Caenorhabditis elegans* during starvation. *Genes & Development*, 21(17), 2161-2171. doi:10.1101/gad.1573107
- Kobet, R. A., Pan, X., Zhang, B., Pak, S. C., Asch, A. S., & Lee, M. (2014). *Caenorhabditis elegans*: a model system for anti-cancer drug discovery and therapeutic target identification. *Biomolecules & Therapeutics*, 22(5), 371-383. doi:10.4062/biomolther.2014.084
- Kyriakakis, E., Markaki, M., & Tavernarakis, N. (2014). *Caenorhabditis elegans* as a model for cancer research. *Molecular & Cellular Oncology*, 2(2). doi:10.4161/23723556.2014.975027
- Lans, H., & Vermeulen, W. (2011). Nucleotide Excision Repair in *Caenorhabditis elegans*. *Molecular Biology International*, 2011, 1-12. doi:10.4061/2011/54279
- Perluigi, M., Domenico, F. D., Blarzino, C., Foppoli, C., Cini, C., Giorgi, A., . . . Coccia, R. (2010). Effects of UVB-induced oxidative stress on protein expression and specific protein oxidation in normal human epithelial keratinocytes: A proteomic approach. *Proteome Science*, 8(1), 13. doi:10.1186/1477-5956-8-13
- Ravindran, J., Prasad, S., & Aggarwal, B. B. (2009). Curcumin and cancer cells: how many ways can curry kill tumor cells selectively? *The AAPS Journal*, 11(3), 495-510. doi:10.1208/s12248-009-9128-x
- Shakeri, A., Cicero, A. F., Panahi, Y., Mohajeri, M., & Sahebkar, A. (2018). Curcumin: a naturally occurring autophagy modulator. *Journal of Cellular Physiology*, 234(5), 5643-5654. doi:10.1002/jcp.27404

Vallianou, N. G., Evangelopoulos A., Schizas, N., & Kazazis, C. (2015). Potential anticancer properties and mechanisms of action of curcumin. *Anticancer Research*, 35(2), 645-651. Retrieved July 28, 2019 from <http://ar.iiarjournals.org/content/35/2/645.full.pdf+html>

Wawruszak, A., Czerwonka, A., Okła, K., & Rzeski, W. (2015). Anticancer effects of ethanol *Lycium barbarum* (goji berry) extract on human breast cancer T47D cell line. *Natural Product Research*, 30(17), 1993-1996. doi:10.1080/14786419.2015.1101691

Zhou, S., Cheng, J., Zhou, Z., Sheng, H., He, L., Fan, X., . . . Zhao, R. J. (2014). An evidence-based update on the pharmacological activities and possible molecular targets of *Lycium barbarum* polysaccharides. *Drug Design, Development and Therapy*, 9, 33-78. doi:10.2147/dddt.s72892

Zhou, S., Zhang, S., Shen, H., Chen, W., Xu, H., Chen, X., . . . Tang, J. (2017). Curcumin inhibits cancer progression through regulating expression of microRNAs. *Tumor Biology*, 39(2), 1-12. doi:10.1177/1010428317691680